

DERMATITIS, CONTACT (ACUTE)

Level I

(If subjective data is sufficient, no exam is necessary.)

Skill Level: RN, LPN

Definition: A delayed (Type 4) fixed inflammatory response evoked when skin contact with an irritant or allergen. Takes one to seven days to fully develop and generally takes seven to 14 days to resolve from a single exposure (poison oak, plants, metal...).

<p>Subjective:</p> <ul style="list-style-type: none">• Clear exposure history.<ul style="list-style-type: none">- Signs and symptoms within 7 days of exposure.- Suspected or actual exposure to a common allergen (e.g.: poison oak) or individual allergen (e.g.: soap, metal, lotion).• No breathing difficulties.• No prior history of serious reaction to the same allergen.• Itchy rash.• Area involved.• "I am allergic to (wool, poison oak, etc.)"	<p>Assessment:</p> <ul style="list-style-type: none">• Altered skin integrity: Dermatitis, Contact• Potential for skin infection.
<p>Objective:</p> <ul style="list-style-type: none">• Distribution of rash consistent with exposure to suspected allergen.• Edematous or vesicular tissue on a red base frequently found in a line or patches on body part exposed to allergen.• Vesicles may be full of clear fluid.• No pustules or other sign of infection.• Area of rash should be non-tender.• No wheezing.• Vital signs normal for patient.	<p>Plan:</p> <p><u>Self care canteen product:</u></p> <ul style="list-style-type: none">• Hydrocortisone cream applied directly to affected area BID x 1 wk for itch. <p>At nursing discretion may use any of the below:</p> <ul style="list-style-type: none">• Check for allergies to medications.• Remove irritant or allergen, (<u>may need to launder bedding and clothing</u>).• Apply cold compress, or• If wool allergy is suspected and not documented in the health care record, schedule wool patch test (or may apply wool patch test at the time of this visit).• Consider Triamcinolone 0.1% ointment BID for 2 weeks if the cause of the rash is clear and it seems to be spreading rapidly. (Do not apply to the face).

Dermatitis, Contact (Acute) - Level I

Nursing Education:

1. Herpes skin infections, Herpes Simplex or Zoster, are caused by a herpes virus. This condition can present as vesicles filled with clear fluid but there is usually tenderness or a burning pain associated with these conditions.
2. While Poison Oak is not contagious, the resin can be on clothing and bedding, so these should be laundered after known exposure.

Patient Education:

1. Contact Dermatitis is non-contagious.
2. Advise patient of the signs and symptoms of second bacterial infection.
3. Advise patient that it may take up to two weeks for condition to resolve. If no improvement within one week, patient should notify Health Services.

APPROVED:

Medical Services Manager

Date

Chief Medical Officer

Date

Steve Shepherd MD

Medical Director

10/6/09

Date

Effective Date: 10/09
Revised: September 2009